

Implant After Care

Guidelines for After Care of your Mouth • Take care to only have cold drinks and do not eat until the local anesthetic has worn off. Avoid hot drinks or hot food for the first day and do not 'swill' liquid over the area. Try not to disturb the area with your tongue or fingers. Do not undertake strenuous exercise for the first 48 hours (running / gym). • You may have some swelling and/or bruising following your treatment - this usually reaches a peak 2 to 3 days later. This is quite normal and both will subside naturally after a few days. Swelling can be reduced with ice packs (or a bag of frozen peas) wrapped in a towel. Hold on the cheek area for a maximum of 10 minutes at a time with 20 minutes break. After 24 hours gentle heat is more beneficial. Sleeping propped up slightly on 2-3 pillows may also help. • Pain should not be a big problem. Although you may be sore for a few days after any surgery in your mouth, this can be easily managed with simple analgesics (pain killers). You should take these regularly at the maximum stated dose for the first 2 days after your surgery. Take whatever painkillers you normally take for headaches, aches and sprains (ibuprofen), and take your first dose before the local anesthetic has completely worn off. • If after a few days you experience increasing pain and swelling, you must return to the office as soon as possible so that we can ensure you are not beginning to develop an infection. • If you have a denture that covers the surgical area please wear it as little as possible for the first week to protect the surgical site during its initial healing period. You should always leave the denture out at night. • The stitches are dissolvable but often remain for around 2-3 weeks, if they are uncomfortable or annoying, you may contact us to remove them. • Some minor bleeding after surgery in the mouth is normal. If this persists, apply pressure by biting firmly down over the area on a dampened gauze swab or clean handkerchief for 60 minutes whilst sitting upright. Do not keep checking or changing the gauze. You should contact us if bleeding persists for any reason after applying pressure in this way. • If you have been given a course of antibiotics to take after your surgery, please ensure that you complete the course. Cleaning • Successful oral surgery depends on keeping the mouth as clean as possible. • Avoid the surgical site for the first few days, but then begin to carefully clean this area as well when tenderness permits. at least 1 minute, 3 times a day, for 7 days. • You should also start cleaning your other teeth as normal. • Try to keep food away from the surgical area for as long as possible. Rinse following eating to keep the area clean. • You are advised not to smoke until the wound has healed as this severely limits healing in the mouth. • We want your recovery to be as smooth and pleasant as possible. It is vital to follow these instructions very carefully - if you have any concerns or questions regarding your progress, please do not hesitate to contact us. An out of hours contact number is always available on our answer phone 407-671-1017